

LAMPIRAN

Neuropati

		Frequency	Percent	Valid Percent	Cumulativ e Percent
Valid	Normal	13	17,3	17,3	17,3
	Ringan	37	49,3	49,3	66,7
	Sedang	25	33,3	33,3	100,0
	Total	75	100,0	100,0	

Nefropati

		Frequency	Percent	Valid Percent	Cumulativ e Percent
Valid	Normal	68	90,7	90,7	90,7
	Ringan	5	6,7	6,7	97,3
	Sedang	2	2,7	2,7	100,0
	Total	75	100,0	100,0	

jeniskelamin

		Frequency	Percent	Valid Percent	Cumulativ e Percent
Valid	Laki-laki	31	41,3	41,3	41,3
	Perempuan	44	58,7	58,7	100,0
	Total	75	100,0	100,0	

Neuropati * BMI Crosstabulation

		BMI					Total
		Normal	Kurus	Obese I	Obese II	Beresiko	
Tidak ada	Count	8	1	1	1	2	13
	% within Neuropati	61,5%	7,7%	7,7%	7,7%	15,4%	100,0%
	% of Total	10,7%	1,3%	1,3%	1,3%	2,7%	17,3%
Ringan	Count	13	4	3	4	11	35
	% within Neuropati	37,1%	11,4%	8,6%	11,4%	31,4%	100,0%
	% of Total	17,3%	5,3%	4,0%	5,3%	14,7%	46,7%
Sedang	Count	9	1	9	4	4	27
	% within Neuropati	33,3%	3,7%	33,3%	14,8%	14,8%	100,0%
	% of Total	12,0%	1,3%	12,0%	5,3%	5,3%	36,0%
Total	Count	30	6	13	9	17	75
	% within Neuropati	40,0%	8,0%	17,3%	12,0%	22,7%	100,0%
	% of Total	40,0%	8,0%	17,3%	12,0%	22,7%	100,0%

Neuropati * Hipertensi Crosstabulation

			Hipertensi			Total
			Normal	Pre-hipertensi	Hipertensi 1	
Neuropati	Tidak ada	Count	12	1	0	13
		% within Neuropati	92,3%	7,7%	,0%	100,0%
		% of Total	16,0%	1,3%	,0%	17,3%
	Ringan	Count	26	8	1	35
		% within Neuropati	74,3%	22,9%	2,9%	100,0%
		% of Total	34,7%	10,7%	1,3%	46,7%
	Sedang	Count	5	10	12	27
		% within Neuropati	18,5%	37,0%	44,4%	100,0%
		% of Total	6,7%	13,3%	16,0%	36,0%
Total	Count	43	19	13	75	
	% within Neuropati	57,3%	25,3%	17,3%	100,0%	
	% of Total	57,3%	25,3%	17,3%	100,0%	

Neuropati * Merokok Crosstabulation

			Merokok		Total
			Ya	Tidak	
Neuropati	Tidak ada	Count	0	13	13
		% within Neuropati	,0%	100,0%	100,0%
		% of Total	,0%	17,3%	17,3%
	Ringan	Count	13	22	35
		% within Neuropati	37,1%	62,9%	100,0%
		% of Total	17,3%	29,3%	46,7%
	Sedang	Count	13	14	27
		% within Neuropati	48,1%	51,9%	100,0%
		% of Total	17,3%	18,7%	36,0%
Total	Count	26	49	75	
	% within Neuropati	34,7%	65,3%	100,0%	
	% of Total	34,7%	65,3%	100,0%	

Proteinuria * BMI Crosstabulation

		BMI					Total
		Normal	Kurus	Obese I	Obese II	Beresiko	
Normal	Count	28	6	11	6	17	68
	% within Proteinuria	41,2%	8,8%	16,2%	8,8%	25,0%	100,0%
	% of Total	37,3%	8,0%	14,7%	8,0%	22,7%	90,7%
Ringan	Count	1	0	2	2	0	5
	% within Proteinuria	20,0%	,0%	40,0%	40,0%	,0%	100,0%
	% of Total	1,3%	,0%	2,7%	2,7%	,0%	6,7%
Sedang	Count	1	0	0	1	0	2
	% within Proteinuria	50,0%	,0%	,0%	50,0%	,0%	100,0%
	% of Total	1,3%	,0%	,0%	1,3%	,0%	2,7%
Total	Count	30	6	13	9	17	75
	% within Proteinuria	40,0%	8,0%	17,3%	12,0%	22,7%	100,0%

Proteinuria * BMI Crosstabulation

		BMI					Total
		Normal	Kurus	Obese I	Obese II	Beresiko	
Normal	Count	28	6	11	6	17	68
	% within Proteinuria	41,2%	8,8%	16,2%	8,8%	25,0%	100,0%
	% of Total	37,3%	8,0%	14,7%	8,0%	22,7%	90,7%
Ringan	Count	1	0	2	2	0	5
	% within Proteinuria	20,0%	,0%	40,0%	40,0%	,0%	100,0%
	% of Total	1,3%	,0%	2,7%	2,7%	,0%	6,7%
Sedang	Count	1	0	0	1	0	2
	% within Proteinuria	50,0%	,0%	,0%	50,0%	,0%	100,0%
	% of Total	1,3%	,0%	,0%	1,3%	,0%	2,7%
Total	Count	30	6	13	9	17	75
	% within Proteinuria	40,0%	8,0%	17,3%	12,0%	22,7%	100,0%
	% of Total	40,0%	8,0%	17,3%	12,0%	22,7%	100,0%

Proteinuria * Hipertensi Crosstabulation

			Hipertensi			Total
			Normal	Pre-hipertensi	Hipertensi 1	
Proteinuria	Normal	Count	41	16	11	68
		% within Proteinuria	60,3%	23,5%	16,2%	100,0%
		% of Total	54,7%	21,3%	14,7%	90,7%
	Ringan	Count	2	2	1	5
		% within Proteinuria	40,0%	40,0%	20,0%	100,0%
		% of Total	2,7%	2,7%	1,3%	6,7%
	Sedang	Count	0	1	1	2
		% within Proteinuria	,0%	50,0%	50,0%	100,0%
		% of Total	,0%	1,3%	1,3%	2,7%
Total	Count	43	19	13	75	
	% within Proteinuria	57,3%	25,3%	17,3%	100,0%	
	% of Total	57,3%	25,3%	17,3%	100,0%	

Proteinuria * Merokok Crosstabulation

			Merokok		Total
			Ya	Tidak	
Proteinuria	Normal	Count	23	45	68
		% within Proteinuria	33,8%	66,2%	100,0%
		% of Total	30,7%	60,0%	90,7%
	Ringan	Count	2	3	5
		% within Proteinuria	40,0%	60,0%	100,0%
		% of Total	2,7%	4,0%	6,7%
	Sedang	Count	1	1	2
		% within Proteinuria	50,0%	50,0%	100,0%
		% of Total	1,3%	1,3%	2,7%
Total	Count	26	49	75	
	% within Proteinuria	34,7%	65,3%	100,0%	
	% of Total	34,7%	65,3%	100,0%	

Correlations

			neuropati	proeinuria
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**
		Sig. (2-tailed)	.	.006
		N	75	75
	proeinuria	Correlation Coefficient	.317**	1.000
		Sig. (2-tailed)	.006	.
		N	75	75

** . Correlation is significant at the 0.01 level (2-tailed).

P=0,006

Uji table kedua klinis dan jenis kelamin (**p=0,625**)

Correlations

			neuropati	proeinuria	jeniskelamin
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**	-.147
		Sig. (2-tailed)	.	.006	.207
		N	75	75	75
	proeinuria	Correlation Coefficient	.317**	1.000	-.057
		Sig. (2-tailed)	.006	.	.625
		N	75	75	75
	jeniskelamin	Correlation Coefficient	-.147	-.057	1.000
		Sig. (2-tailed)	.207	.625	.
		N	75	75	75

Correlations

			neuropati	proeinuria	jeniskelamin
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**	-.147
		Sig. (2-tailed)	.	.006	.207
		N	75	75	75
	proeinuria	Correlation Coefficient	.317**	1.000	-.057
		Sig. (2-tailed)	.006	.	.625
		N	75	75	75
	jeniskelamin	Correlation Coefficient	-.147	-.057	1.000
		Sig. (2-tailed)	.207	.625	.
		N	75	75	75

** . Correlation is significant at the 0.01 level (2-tailed).

Tabel ketiga BMI (p=0,067)

Correlations

			neuropati	proeinuria	bmi
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**	.241
		Sig. (2-tailed)	.	.006	.038
		N	75	75	75
	proeinuria	Correlation Coefficient	.317**	1.000	.213
		Sig. (2-tailed)	.006	.	.067
		N	75	75	75
	bmi	Correlation Coefficient	.241	.213	1.000
		Sig. (2-tailed)	.038	.067	.
		N	75	75	75

** . Correlation is significant at the 0.01 level (2-tailed).

Correlations

			neuropati	proeinuria	bmi
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**	.241
		Sig. (2-tailed)	.	.006	.038
		N	75	75	75
	proeinuria	Correlation Coefficient	.317**	1.000	.213
		Sig. (2-tailed)	.006	.	.067
		N	75	75	75
	bmi	Correlation Coefficient	.241	.213	1.000
		Sig. (2-tailed)	.038	.067	.
		N	75	75	75

*. Correlation is significant at the 0.05 level (2-tailed).

Tabel tekanan darah (p=0,054)

Correlations

			proeinuria	TD
Spearman's rho	proeinuria	Correlation Coefficient	1.000	.223
		Sig. (2-tailed)	.	.054
		N	75	75
	TD	Correlation Coefficient	.223	1.000
		Sig. (2-tailed)	.054	.
		N	75	75

Tabel status merokok (p=0,631)

Correlations

			proeinuria	merokok
Spearman's rho	proeinuria	Correlation Coefficient	1.000	-.056
		Sig. (2-tailed)	.	.631
		N	75	75
	merokok	Correlation Coefficient	-.056	1.000
		Sig. (2-tailed)	.631	.
		N	75	75

Descriptives^a

Neuropati			Statistic	Std. Error		
Nefropati	Ringan	Mean	,05	,038		
		95% Confidence Interval for Mean				
		Lower Bound	-,02			
		Upper Bound	,13			
		5% Trimmed Mean	,00			
		Median	,00			
		Variance	,053			
		Std. Deviation	,229			
		Minimum	0			
		Maximum	1			
		Range	1			
		Interquartile Range	0			
		Skewness	4,113	,388		
		Kurtosis	15,767	,759		
			Sedang	Mean	,28	,123
				95% Confidence Interval for Mean		
Lower Bound	,03					
Upper Bound	,53					
5% Trimmed Mean	,20					
Median	,00					
Variance	,377					
Std. Deviation	,614					
Minimum	0					
Maximum	2					
Range	2					
Interquartile Range	0					
Skewness	2,127			,464		
Kurtosis	3,539			,902		

a. Nefropati is constant when Neuropati = Normal. It has been omitted.

Tests of Normality^b

Neuropati		Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Nefropati	Ringan	,539	37	,000	,241	37	,000
	Sedang	,476	25	,000	,515	25	,000

a. Lilliefors Significance Correction

b. Nefropati is constant when Neuropati = Normal. It has been omitted.

Correlations

			neuropati	proeinuria
Spearman's rho	neuropati	Correlation Coefficient	1.000	.317**
		Sig. (2-tailed)	.	.006
		N	75	75
	proeinuria	Correlation Coefficient	.317**	1.000
		Sig. (2-tailed)	.006	.
		N	75	75

** . Correlation is significant at the 0.01 level (2-tailed).

Test of Homogeneity of Variance^a

		Levene Statistic	df 1	df 2	Sig.
Nefropati	Based on Mean	19,359	1	60	,000
	Based on Median	4,180	1	60	,045
	Based on Median and with adjusted df	4,180	1	34,101	,049
	Based on trimmed mean	14,828	1	60	,000

a. Nefropati is constant when Neuropati = Normal. It has been omitted.

Nefropati * Neuropati Crosstabulation

Count

		Neuropati			Total
		Normal	Ringan	Sedang	
Nefropati	Normal	13	35	20	68
	Ringan	0	2	3	5
	Sedang	0	0	2	2
Total		13	37	25	75