

**L
A
M
P
I
R
A
N**

Lampiran IPA

A. Case Summary IPA

Case Processing Summary

Tahfidz	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Nilai_UN kurang dari 4	56	100.0%	0	.0%	56	100.0%
lebih dari atau sama dengan 4	37	100.0%	0	.0%	37	100.0%

B. Deskriptife data

Descriptives

Tahfidz		Statistic	Std. Error	
Nilai_UN kurang dari 4	Mean	2.3688E2	6.11611	
	95% Confidence Interval for Mean	Lower Bound	2.2462E2	
		Upper Bound	2.4913E2	
	5% Trimmed Mean	2.3610E2		
	Median	2.3225E2		
	Variance	2.095E3		
	Std. Deviation	4.57688E1		
	Minimum	144.50		
	Maximum	370.00		
	Range	225.50		
	Interquartile Range	59.50		
	Skewness	.230	.319	
	Kurtosis	.310	.628	
	lebih dari atau sama dengan 4	Mean	2.5800E2	6.74979
95% Confidence Interval for Mean		Lower Bound	2.4431E2	
		Upper Bound	2.7169E2	
5% Trimmed Mean		2.5935E2		
Median		2.5850E2		
Variance		1.686E3		
Std. Deviation		4.10574E1		
Minimum		171.00		
Maximum		320.50		
Range		149.50		
Interquartile Range		64.25		
Skewness		-.316	.388	
Kurtosis		-.802	.759	

C. Tes Normalitas IPA

Tests of Normality

Tahfidz	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	Df	Sig.	Statistic	Df	Sig.
Nilai_UN kurang dari 4	.066	56	.200*	.985	56	.699
lebih dari atau sama dengan 4	.130	37	.117	.956	37	.153

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Karena > 50 menggunakan komolgrov smirnov
(Normal karena $> 0,05$)

D. Tes Homogenitas IPA

Test of Homogeneity of Variance

	Levene Statistic	df1	df2	Sig.
Nilai_UN Based on Mean	.007	1	91	.933
Based on Median	.002	1	91	.965
Based on Median and with adjusted df	.002	1	83.249	.965
Based on trimmed mean	.005	1	91	.943

E. Hasil independen T tes

Group Statistics

tahfidz	N	Mean	Std. Deviation	Std. Error Mean
Nilai_UN kurang dari 4	56	2.3688E2	45.76881	6.11611
lebih dari atau sama dengan 4	37	2.5800E2	41.05738	6.74979

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	T	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Nilai_UN	.007	.933	-2.268	91	.026	-21.12500	9.31445	-39.62701	-2.62299
			-2.319	82.834	.023	-21.12500	9.10860	-39.24217	-3.00783

F. Tes korelasi

Correlations

		tahfidz	Nilai_UN
tahfidz	Pearson Correlation	1	.231*
	Sig. (2-tailed)		.026
	N	93	93
Nilai_UN	Pearson Correlation	.231*	1
	Sig. (2-tailed)	.026	
	N	93	93

*. Correlation is significant at the 0.05 level (2-tailed).

Lampiran IPS

A. Frekuensi data seluruhnya

Statistics

Tahfidz

N	Valid	141
	Missing	0
Mean		3.8901
Std. Error of Mean		.34246
Median		2.9615 ^a
Mode		3.00
Std. Deviation		4.06645
Variance		16.536
Range		29.00
Minimum		1.00
Maximum		30.00
Sum		548.50
Percentiles	10	1.4176 ^b
	20	1.8432
	25	2.0338
	30	2.2243
	40	2.6000
	50	2.9615
	60	3.3150
	70	3.8722
	75	4.3167
	80	4.8071
	90	6.3111

a. Calculated from grouped data.

b. Percentiles are calculated from grouped data.

Frekuensi data IPS

Statistics

		Tahfidz	Nilai_UN
N	Valid	63	48
	Missing	78	93
Mean		1.22	2.4975E2
Std. Error of Mean		.053	6.92843
Median		1.22 ^a	2.4825E2 ^a
Mode		1	240.50
Std. Deviation		.419	4.80015E1
Variance		.176	2.304E3
Range		1	211.50
Minimum		1	161.50
Maximum		2	373.00
Sum		77	1.20E4
Percentiles	10	. ^{b,c}	1.9040E2 ^c
	20	.	2.1060E2
	25	.	2.1300E2
	30	.	2.1625E2
	40	1.02	2.3670E2
	50	1.22	2.4825E2
	60	1.42	2.6230E2
	70	1.62	2.7315E2
	75	1.72	2.7675E2
	80	1.82	2.8140E2
90	.	3.1892E2	

a. Calculated from grouped data.

D. Case proesing summary IPS

Case Processing Summary

Tahfidz	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Nilai_UN kurang dari 4	38	77.6%	11	22.4%	49	100.0%
lebih dari atau sama dengan 4	10	71.4%	4	28.6%	14	100.0%

E. Tes Normalitas IPS

Tests of Normality

Tahfidz	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	Df	Sig.
Nilai_UN kurang dari 4	.118	38	.200*	.945	38	.061
lebih dari atau sama dengan 4	.187	10	.200*	.954	10	.713

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Karena jumlah data < 50 maka menggunakan normalitas saphiro wilk (Normal karena > 0,05)

F. Tes homogenitas IPS

Test of Homogeneity of Variance

	Levene Statistic	df1	df2	Sig.
Nilai_UN Based on Mean	4.110	1	46	.048
Based on Median	3.877	1	46	.055
Based on Median and with adjusted df	3.877	1	41.795	.056
Based on trimmed mean	3.881	1	46	.055

G. Hasil Independent sample T test

Group Statistics

Tahfidz	N	Mean	Std. Deviation	Std. Error Mean
Nilai_UN kurang dari 4	38	2.4266E2	50.20769	8.14476
lebih dari atau sama dengan 4	10	2.7669E2	25.50806	8.06636

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Nilai- Equal UN variances assumed	4.110	.048	-2.063	46	.045	-34.03211	16.49847	-67.24181	-.82241
Equal variances not assumed			-2.969	29.299	.006	-34.03211	11.46313	-57.46645	-10.59776

H. Tes korelasi

Correlations

		Tahfidz	Nilai_UN
Tahfidz	Pearson	1	.291*
	Correlation		
	Sig. (2-tailed)		.045
	N	63	48
Nilai_UN	Pearson	.291*	1
	Correlation		
	Sig. (2-tailed)	.045	
	N	48	48