

DAFTAR PUSTAKA

- Amerongen. (1991). *Ludah dan Kelenjar Ludah Arti Bagi Kesehatan Gigi*. Yogyakarta: Gadjah Mada University Press.
- Barrett, K. E., Boitano, S., Barman, S. M., & Brooks, H. L. (2012). Overview. Gastrointestinal Physiology. In *Ganong's Review of Medical Physiology, 24e*.
- Bhalajhi, S. I. (2004). *Orthodontics the art and science 3 rd ed*. New Delhi: Arya (MEDI) Publishing House. h. 271, 384.
- Bonetti, G. A., Incerti Parenti, S., Garulli, G., Gatto, M. R., & Checchi, L. (2013). Effect of fixed orthodontic appliances on salivary properties. *Progress in Orthodontics, 14*, 13.
- Carillo EL, Bastida NMM, Perez LS, Tavora JA. (2010). Effect of orthodontic treatment on saliva, plaque and the levels of streptococcus and lactobacillus. *Med Oral Patol Oral Cir Bucal, 15*(6): 924-929.
- Carillo EL, Bastida NMM, Perez LS, Tavora JA. (2010). Changes in the oral environment during four stages of orthodontic treatment. *Korean J Orthod, 40*(2): 95-105.
- Carlson, E. R., & Ord, R. A. (2015). *Salivary Gland Pathology: Diagnosis and Management. Salivary Gland Pathology: Diagnosis and Management*.
- Chang, H. S., Walsh, L. J., Freer, T. J., & Chang HS, Walsh LJ, F. T. (1999). The effect of orthodontic treatment on salivary flow, pH, buffer capacity, and levels of mutans streptococci and lactobacilli. *Australian Orthodontic Journal, 15*, 229–34.
- Cunningham, F. G., Gant N. F., Leveno K. J., Gilstrap III L. C., Hauth J. C., Wenstrom, K. D. (2006). *Fisiologi Kehamilan*. In: Hartanto Huriawati, dkk. (eds.) *Obstetri Williams*. 21st ed. Jakarta: EGC. p180-213.
- Degerman, A., Chun, D., Nguyen, T. B., Bravo, D. T., Alanis, J., Rokaeus, A., & Waschek, J. A. (2002). Local action of estrogen and thyroid hormone on vasoactive intestinal peptide (VIP) and galanin gene expression in the rat anterior pituitary. *Neuropeptides, 36*(1), 50–57.
- Finn, S. B. (2003). *Clinical Pedodontics 4thed*, WB. Philadelphia: Saunders Co.
- Flink, H. (2007). Studies on the prevalence of reduced salivary flow rate in relation to general health and dental caries, and effect of iron supplementation. *Swedish dental journal. Supplement*.
- Foster, T.D., 1997. *Buku Ajar Ortodontisi*. Translated by L. Yuwono. Jakarta: EGC.

- Gavião, M. B. D., & Bilt, A. Van Der. (2004). Salivary secretion and chewing: stimulatory effects from artificial and natural foods. *Journal of Applied Oral Science : Revista FOB*, 12(2), 159–63.
- Guyton, A. C., & Hall, J. E. (2011). *Guyton Dan Hall Buku Ajar Fisiologi Kedokteran* (Vol. XXXIII).
- Höld, K. M., Boer, D. De, Ph, D., Zuidema, J., & Maes, R. a a. (2012). Saliva As an Analytical Tool Saliva As an Analytical Tool in Toxicology. *Utrecht Institute of Pharmaceutical Science (UIPS)*, 1(1), 1–34.
- Indriana, T. (2011). Perbedaan Laju Aliran Saliva dan pH karena Pengaruh Stimulus Kimiawi dan Mekanis. *J. Kedokt Meditek*, 17(44), 1–5.
- Isaacson, K., J-D-Muir, & Reed, R. (2002). *Removable Orthodontics Appliances*. New Delhi: Elsevier.
- Joskow, J. and Yamane, T. (1965) ‘Statistics, an Introductory Analysis.’, *Journal of the American Statistical Association*, 60(310), p. 678. Kasuma N, Biomod M. (2015). *Fisiologi dan Patologi Saliva*. Padang. h.22-5
- Kidd, Edwina A.M, Sally Joyston-Bechal. (2012). *Dasar-dasar Karies Penyakit dan Penanggulangannya*. Jakarta: EGC. h. 145-52.
- Kuswandani, F. (2016). Analisis Kadar Kalsium Saliva dan Hubungannya dengan Pembentukan Karang Gigi. *Indonesian Journal of Pharmaceutical Science and Technology*, 3(1), 31-37.
- Modéer, T., Blomberg, C. C., Wondimu, B., Julihn, A., & Marcus, C. (2010). Association between obesity, flow rate of whole saliva, and dental caries in adolescents. *Obesity (Silver Spring, Md.)*, 18(12), 2367–2373.
- Navazesh, M. and Kumar, S. K. S. (2008). Measuring salivary flow: challenges and opportunities. *Journal of the American Dental Association (1939)*. American Dental Association, 139 Suppl(May), p. 35S–40S.
- Peros, K., Mestrovic, S., Anic-Milosevic, S., & Slaj, M. (2011). Salivary microbial and nonmicrobial parameters in children with fixed orthodontic appliances. *Angle Orthodontist*, 81(5), 901–906.
- Rashkova, M. (2012). Influence of Systemic Diseases and Removable Orthodontic Appliances on the Quality of Saliva in Childhood. *Journal of IMAB - Annual Proceeding (Scientific Papers)*, 18, 2(2012), 163–167.
- Rhoades, R., & Bell, D. (2009). *Medical Physiology: Principles for Clinical Medicine*. Lippincott Williams & Wilkins.
- Rockenbach, M. I., Marinho, S. A., Veeck, E. B., Lindemann, L., & Shinkai, R. S. (2006). Salivary flow rate, pH, and concentrations of calcium, phosphate, and sIgA in Brazilian pregnant and non-pregnant women. *Head & Face Medicine*, 2(44), 1–5.

- Saloom, H. F., Mohammed-Salih, H. S. and Rasheed, S. F. (2013). The influence of different types of fixed orthodontic appliance on the growth and adherence of microorganisms (in vitro study). *Journal of Clinical and Experimental Dentistry*, 5(1), pp. 36–41.
- Sherwood, L. (2010). Human Physiology: From Cells to Systems. *Human Physiology, 7th edition*, 766.
- Singh, G. (2007). *Textbook of orthodontics*. 2nd edn, Jaypee, India. 2nd edn. doi: 10.1017/CBO9781107415324.004.
- Soesilo, D., Santoso, R. E., & Diyatri, I. (2005). Peranan sorbitol dalam mempertahankan kestabilan pH saliva pada proses pencegahan karies. *Majalah Kedokteran Gigi (Dent. J)*, 38(1), 25–28.
- Sulandjari, H. (2008). *Buku Ajar Orthodonti 1*. Yogyakarta: Fakultas Kedokteran Gigi UGM.
- Takeuchi, K., Furuta, M., Takeshita, T., Shibata, Y., Shimazaki, Y., Akifusa, S., Yamashita, Y. (2015). Risk factors for reduced salivary flow rate in a Japanese population: The Hisayama study. *BioMed Research International*, 2015.
- Välilä, H., Savolainen, S., Soukka, T., Silvoniemi, P., Mäkelä, S., Kujari, H., Laine, M. (2004). Estrogen receptor- is the predominant estrogen receptor subtype in human oral epithelium and salivary glands. *The Journal of Endocrinology*, 180(1), 55–62.