

The Effect of Ethanolic Extract of Purple Yam Tuber (*Dioscorea alata* L.) on Bone Calcium Levels in Ovariectomized Rat

by Nurul Makiyah

Submission date: 25-Sep-2019 10:40AM (UTC+0700)

Submission ID: 1179553719

File name: 2._Hasil_Review_Journal_of_Health_Medicine_Nursing_Vol_17.pdf (157.33K)

Word count: 3259

Character count: 17484

The Effect of Ethanolic Extract of Purple Yam Tuber (*Dioscorea alata* L.) on Bone Calcium Levels in Ovariectomized Rat

Sri Tasminatun

Pharmacology Department,

The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta

AlfainaWahyuni

Obstetrics and Gynecology Sciences Department,

The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta

S.N. NurulMakiyah

Histology Department

The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta

Zulfianisa

Pharmacy Student

The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta

Abstract

This study is on the effects of the ethanolic extract of purple yam tuber (*Dioscoreaalata* L.) on the bone calcium levels in ovariectomized (ovx) rat models of osteoporosis. Forty-five of 8-weeks-old female Sprague–Dawley rats were randomly assigned to six groups as followed: sham-operated, OVX, OVX-Estradiol, OVX-EDA 116 mg/kg BW, OVX-EDA 232 mg/kg BW, OVX-EDA 463 mg/kg BW for 30 days. The administration of EDA and estradiol was given orally using a stomach tube. It is on the 31st day since the bones of femur and Tibia rat were isolated. The isolated bones were analyzed using wet destruction and the bone calcium level was measured using the AAS. The bone calcium level were tested statistically using one way ANOVA. The average of bone calcium levels for each group: for the normal group is $18,30 \pm 3,17\%$, OVX $16,44 \pm 3,93\%$, OVX-Estradiol $16,59 \pm 2,63\%$, and for the group of OVX-EDA dose 116; 232; 463 are $15,89 \pm 1,66\%$, $15,96 \pm 2,61\%$, and $14,91 \pm 2,31\%$ respectively. The conclusion is that the treatment of purple yam tuber (*Dioscoreaalata*L.) with the different dose of 116; 232; and 463 mg/kg BW for 30 days cannot increase the bone calcium level in the ovariectomized rats. According to the statistical data there is no significant differences among the treatment groups.

Keywords: Calcium, *Dioscorea alata*, Extract

INTRODUCTION

Osteoporosis is a condition where the bones are weak and it is breakable. Osteoporosis is one of the bone metabolic diseases signaled by the decreasing of the bone mass because of the deficit of the bone matrix and mineral along with the damage of the microarchitecture of the bone tissue that causes the decreasing of the bone strength and the risk of bone fracture. ¹National Osteoporosis Foundation states that 44 millions American have the risk to get osteoporosis, and 22% of them are adults up to senior people who mostly are menopause women. ²

Menopause is a condition when the ovariums are unable to produce estrogen. Estrogen hormone has contribution to help the calcium absorption in the intestine to prevent the bone density. The decreasing of the calcium absorption will disrupt the balance of the calcium in the blood because the low blood calcium may lead to the increasing of the calcium reabsorption in the bone that causes low bone mass. The decreasing of the calcium level in the bone because of the estrogen deficiency is called *postmenopausal osteoporosis* or *osteoporosis tipe*.³

The injection of the hormone to substitute the estrogen (*Estrogen Replacement Therapy*-ERT) is commonly given to the menopause women to overcome the symptom that occur in pasca-menopause and to prevent chronic disease like osteoporosis.⁴ The sintetic estrogen preparat given as the therapy to substitute the hormone is expensive as well as it has bad side effects. Syntetic estrogen can interfere the blood coagulation process, disturb enzyme works in the liver, lead to endometrium uterus bleeding, nausea and throw up. ⁵

The use of syntetic estrogen preparate continuously can cause carcinoma ovarium.⁶ Besides, the use of estrogen and progesterone preparate as the hormone replacement therapy can also lead to the breast cancer. ⁷

There are side effects in the hormonal therapy. It is wiser to use phytoestrogen therapy as the alternative for the menopause women because it is saver.⁸ Phytoestrogen is a natural substance that can influence the body

estrogenic activities.⁹

One of the plants that has phytoestrogen substance is the purple yam tuber (*Dioscorea alata* L.). The D tuber has 5 substances that has the estrogenic quality based on the *ligand-dependent*¹⁰ transcription activities. The consumption of *D. alata tuber* can increase the strength of bones in the ovariectomized rats.¹¹

Testing the effect of ethanolic extract of *D. alata* on the calcium level of ovariectomized rats was conducted in this research and measured by using AAS. This research is hoped to be able to prove the effect of ethanolic extract of *D. alata* tuber on the increasing bone calcium level in the ovariectomized rats. Therefore, the tuber is useful to be used as the phytoestrogen agent to prevent the osteoporosis in the menopause women. The result of the research is also useful as the reference in the utilization as well as for other researcher in the pharmacology division. This research aims to know the effect of ethanolic effect of *D. alata* tuber on the bone calcium level in the ovariectomized rats.

MATERIALS AND METHODS

Plant and Chemical Materials

Purple yam tuber (*Dioscoreaalata* L.) *D. alata* used in the present study were collected from the commercial market in Kulonprogo Yogyakarta and authenticated at the Laboratory of Pharmacognosy, The Faculty of Pharmacy, Gadjah Mada University, Yogyakarta, Indonesia. Concentrated citric acid solution, Perchlorate acid solution, distilled water and Calcium Carbonate were obtained from LPPT Gadjah Mada University, Yogyakarta, Indonesia

The Preparation of Ethanolic Extract of Purple Yam Tuber (Dioscoreaalata L.)

The dried powders of tuber *Dioscoreaalata* L were extracted by maceration using ethanolic 70 %. The ethanolic extracts were evaporated to obtain the concentrated extract.

Animals

Female Sprague–Dawley rats, aged 6 week, were purchased from Faculty of Medicine, Gadjah Mada University, Yogyakarta, Indonesia. The animals were grouped and housed in polyacrylic cages with not more than five animals per cage and maintained under the standard laboratory conditions (temperature $25 \pm 2^{\circ}\text{C}$) with dark and light cycle (12/12 h) and allowed free access to commercial pellet diet and water ad libitum.

Administration Procedure

Rats were acclimatized to laboratory condition for 1 week before the commencement of the experiment. Ovariectomized were done to all experimental animals except to the normal group. Animals were assigned to the experimental groups, normal (sham-operated), OVX, OVX-Estradiol, OVX-EDA 116 mg/kg BW, OVX-EDA 232 mg/kg BW, OVX-EDA 463 mg/kg BW, with five animals per group, per experiment. Twenty days after ovariectomy, all of the rats were allowed to the controlled access to a commercial standard pellet and free access to deionized water for 20 days.

The extract of *D. alata* tuber was given orally once a day during 30 days using the sonde technique and the doses are 116 mg/kg BW, 232 mg/kg BW, and 463 mg/kg BW to the experimental group of OVX-EDA. The group of OVX-Estradiol was given estradiol with the dose of 2 $\mu\text{g}/\text{day}$, and the control groups and OVX did not get the treatment. On the 31st day, the rats were sacrificed to take their thigh bones.

The Measurement of Bone Calcium Level Using the Atomic Absorption Spectroscopy (AAS)

The isolated bones then were weighted to find out the weight of the bones. The sample of the isolated bones that had been weighted was to be destructed by using 15 ml HNO_3 and 3 ml HClO_4 , and heated until the bones dissolve in the solution and the solution became clear. The destruction was continued until ± 3 ml left, then added 25 ml of distilled water to it. The solution of the sample was filtered and diluted using the distilled water until 100 ml. The sample as the result of destruction process was ready to be read at the absorbance part using the AAS at the wavelength of (λ) 422,7 nm. The result of the reading then was compared with the standard curve, therefore came up the level of bone calcium in mg/dl or ppm.

The standard curve is the result of the absorbansi measurement from several series dilution from calcium standard dilution. The concentrations of carbonat calcium (CaCO_3) that was used as the standar solution are 0; 0,05; 0,1; 0,2; 0,5; 0,8; 1; 2; 3; 4; and 5 mg/L. The result of standar solution absorbance measurement would create the similarity on the linear lines that were used to find out the calcium level of the ovariectomized rat bones that were used as the sample,

Data Analysis

The normality of the bone calcium level data were tested using the *Saphiro-Wilk*. The results of the normality and varian test data were analyzed using *One Way* ANOVA test, and the measurement was continued using the Tukey test to know the meaningful differences among the treatment groups.

RESULT AND FINDING

The average of bone calcium level was gained from the measurement of bone calcium level in white rats (*Rattus norvegicus*) type *Sprague-Dowley* ovariectomy model. Those rats were given the therapy of ethanolic extract of purple yam tuber *Dioscorea alata* L) and measured by AAS. The result is shown in Table 1.

Table 1. The average of bone calcium level.

No.	Group Treatment	The Average of Bone Ca (%) Level
1.	Normal	18,30 ± 3,17
2.	OVX	16,45 ± 3,94
3.	OVX-Estradiol	16,59 ± 2,63
4.	OVX-EDA dosis 116 mg/kgBW/day	15,89 ± 1,66
5.	OVX-EDA dosis 232 mg/kgBW/day	15,96 ± 2,61
6.	OVX-EDA dosis 463 mg/kgBW/day	14,91 ± 2,31

Ovariectomy is a process to take away the ovary in order to make the model rats become menopause. The assumption is that after the ovariectomy, the rats will experience the decreasing of estrogen hormone and it is similar to the hormone deficiency in menopause women. The decreasing of the estrogen hormone after the ovariectomy because the ovary as the hormone producer is already been removed. The decreasing of estrogen hormone in the animal model is proven by Hartiningsih *et al.*, (2010)¹² who explains that the ovariectomy in rats can decrease drastically the estrogen hormone level, so the condition is similar to menopause.

The bone calcium level is different between normal group and OVX group. The normal rat treatment group is the groups that have normal ovary and they still produce estrogen endogenous hormone. The average of bone calcium level at the normal group is 18.30%, and it is higher than OVX group, 16.45%. Although the difference is not significant based on the statistical analysis. The decreasing of bone calcium in the OVX group is to prove that that estrogen deficiency in the ovariectomy rats also decrease the bone calcium level. Nurrochmad *et al.*, (2010)² also state the same result.

Estrogen endogenous deficiency in the menopause women can lead to the increasing of osteoclastogenesis process and the loss of bone mass. In order to keep the estrogen balance, menopause women need to get substitute for the estrogen hormone. Estrogen intake can increase the apoptosis of osteoclasts as it hinders the formation of cytokines that can stimulate the osteoclasts cell differentiation (IL-1, IL-6, dan TNF- α). Besides, the estrogen intake can also stimulate the *osteoprotegerin* (OPG) expression and TGF- β (*Transforming Growth Factor- β*) from the osteoblast and stroma cells, so the process can hamper bone resorption.^{13,14} It is in line with the result of calcium level in OVX-Estradiol, which get the Estradiol therapy of 2 μ g/day. The average of the calcium level is 16.59% and it is higher than the OVX group.

The mechanism of bone resorption which is mediated by osteoclast cell will damage the bone mineral crystals ligament that have the form of *calcium phosphatase* with the collagen matrix. The interaction of osteoclast with the bone surface during the process of resorption not only release some acids (for example citric acid, lactic acid) from the mitochondria and secretory vesicles, but also release the hydrogen ions through *carbonic anhydrase* dan *vacuolar ATPase pompa proton* channel. This condition creates an acidic environment at the *resorptive pit* that causes damage to the bone matrix.¹⁵ Osteoblast cells have a role to secrete the bone matrix component like collagen and proteoglycan. Collagen has function to give strength to matrix and binding the bone mineral salt. Proteoglycan has function to deposit the bone mineral.¹⁵

Calcium and phosphate salts are deposited on the surface of the bone maker cells as the non-crystal compound which then are converted into *hydroxyapatite* [$\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$] of the bone major crystal salt. The function of *hydroxyapatite* is to keep the ion and give strength to the bone. Some *hydroxyapatites* stay in the form of non-crystal to facilitate the fast absorption when needed.¹⁵

The averages of bone calcium level from the treatment rat group of OVX-EDA 116; 232; 463 are 15.89%; 15.96%, 14.91% respectively. The averages are smaller than OVX and OVX-Estradiol groups. The intakes of *D. alata* tuber extract for 30 days do not give effect to the significant increasing of the calcium level.

D. alata tuber contains phytoestrogen that is useful for the menopause women. Phytoestrogen in *D. alata* tuber is different from the existing phytoestrogen, namely diosgenin. Diosgenin in *D. alata* tuber is a spirostanol saponin that is composed from hydrophilic sugar and bounded at the hydrophobic steroid aglycone. The effect of estrogenic from diosgenin is used as the major precursor of the synthetic steroid production.¹⁶ It has been proven before that the effect of the work mechanism of the estrogenic diosgenin is almost the same as the estrogen hormone, which is to influence the formation process and bone resorption seen from the result of RANKL/OPG by Zhang *et al.*, (2014).¹⁷

The inconsiderable impact posed by the increasing bone calcium level can be caused by many factors that can influence the bone formation process. One of the factors is the inadequate dose of the ethanolic extract of *D. alata* purple yam tuber. Zhang *et al.* states that the intake of diosgenin dose 96 mg/kg BW for 12 weeks

can give the effect of antiosteoporosis in menopause women. Another research says that the content of diosgenin in 100 g *D. alata* is only 82.39 mg diosgenin.¹⁸ Therefore the intake of ethanolic extract of *D. alata* purple yam tuber with the doses of 116; 232; 436 mg/kg Bw a day does not give effect of antiosteoporosis in increasing the bone calcium level because the doses are inadequate and the time is also short. Bone calcium level is not only influenced by estrogen hormone, but it is also influenced by other factors, such as calcitonin, parathyroid hormone and the intake of vitamin D that have role to manage the osteoclasts and osteoblasts function. The result of ANOVA statistic test shows that the bone calcium levels among the group are not significantly different; it is shown from the significant value of $p > 0.05$ (0,554). Furthermore, the *Tukey* test also shows that significant values of all the samples above are more than 0.05, which means that all of the data on the bone calcium levels are not different actually among the treatment.

CONCLUSION

This research prove that the intake of ethanolic extract of purple yam tuber (*Dioscorea alata* L.) with variety of doses 116; 232; and 463 mg/kg BW during 30days cannot increase the bone calcium level in the ovariectomized rats. Another research should be conducted on the effect of ethanolic extract of purple yam tuber (*Dioscorea alata* L.) on the ovariectomizedrats with higher doses and longer treatment.

ACKNOWLEDGEMENT

The sincere gratitude goes to The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta who provides the research fund through the Regular Research Program of The Faculty of Medical and Health Sciences Universitas Muhammadiyah Yogyakarta year 2014.

REFERENCES

- ¹Kawiyana, (2009).Osteoporosis Patogenesis Diagnosis Dan PenangananTerkini.Sub Bagian/ SMF Orthopaedi&Traumatologi. BagianBedah FK UNUD / RSUP Sanglah Denpasar.*Jurnal Penyakit Dalam, Volume 10 Nomor 2*.
- ²Nurrochmad A., Leviana F., Wulancarsari C. G., Lukitaningsih E., (2010). Phytoestrogens of *Pachyrhizuserosus* Prevent Bone Loss in an Ovariectomized Rat Model of Osteoporosis.*International Journal of Phytomedicine* 2 363-372. ISSN: 0975-0185.
- ³Lane, Nancy E. (1999).*Lebih Lengkap Tentang : OSTEOPOROSIS Rapuh Tulang*, (Eri D. Nasution, Penerjemah). Jakarta: PT Raja Grafindo Persada.
- ⁴Departemen Kesehatan Republik Indonesia.(2008). *Pedoman Pengendalian Osteoporosis. Keputusan Menteri Kesehatan Republik Indonesia* No. 1142/MENKES/SK/XII/2008:3-4.
- ⁵Wratsangka, R., (1999). Pemberian Terapi Sulih Hormon Sebagai Upaya Meningkatkan Kesehatan Wanita Menopouse. 155-162. *J Kedokter Trisakti, September-Desember 1999-Vol.18, No.3*.
- ⁶Gang P, Kerlikowske K, Subak L, dan Grady D. (1999).*Hormone Replacement Therapy and the Risk of Epithelial Ovarian Carcinoma: a Meta-Analysis*. Obs Gyn.92: 472-479.
- ⁷Ross RK, Paganini-Hill A, Wan PC, dan Pike MC., (2000).*Effect of Hormone Replacement Therapy on Breast Cancer Risk: Estrogen Versus Estrogen Plus Progestin*. Natl Cancer Inst.92: 328-32.
- ⁸Rimoldi, G., et al. (2007).Effect of Chronic Genistein Treatment in Mammary Gland, Uterus, and Vagina.*Jurnal, Jerman, Universitas Georg-August*.
- ⁹Biben H.A., (2012).*Fitoestrogen: Khasiat Terhadap Sistem Reproduksi, Non Reproduksi dan Keamanan Penggunaannya*. Disampaikan dalam Seminar Ilmiah Nasional Estrogen Sebagai Sumber Hormon Alam pada 31 maret 2012.
- ¹⁰Cheng Wei-Yi, Kuo Yueh-Hsiung, & Huang Ching-Jang.(2007). Isolation and Identification of Novel Estrogenic Coumpounds in Yam Tuber (*Dioscoreaalata* CV. Tainung No. 2).*Journal of Agricultural and Food Chemistry*. Vol. 55 No. 18, 7350-7358.
- ¹⁰hen Jian-Hong, Wu James, Lin Hsiao-Che, Wu Shey-Lin, Wang Wen-Fu, Huang Shu-Kuei., et al. (2008). *Dioscorea improves the morphometric and mechanical properties of bone in ovariektomised rats*. *Journal of the Science of Food and Agriculture* 88:2700-2706.
- ¹²Hartiningih, Devita, A., Irkham, W., (2010). *Pengaruh Panhisterektomi dan Konsumsi Suplemen 1,25-Dihidroksivitamin D3 Selama 1,5 Bulan terhadap Retensi Kalsium pada Tikus Wistar*. Lembaga Penelitian Universitas Gadjah Mada.
- ¹³National Osteoporosis Foundation.*Clinician's Guide to Prevention and Treatment of Osteoporosis*. Washington DC: National Osteoporosis Foundation (2010).
- ¹⁵Kennedy, P., Jain, A., Srinivasan, R., and Chaudhry, S. (2008). Bone Homeostasis. *Artikel McMaster Pathophysiology Review*. Diakses pada 10 Juni 2015, dari <http://www.pathophys.org/osteoporosis/>
- ¹⁶Prabowo, A.Y., Estiasih, T., Purwatiningrum, I., (2014).Gembili (*Dioscoreaesculata*L.) as Food Contain

Bioactive compound: A Review. *Jurnal Pangandan Agroindustri Vol.2 No.3 p.129-135, Juli 2014.*

¹⁷Zhang, Z., Song, C., Fu, X., Liu, M., *et al.*, (2014). High-Dose Diosgenin Reduces Bone Loss in Ovariectomized Rats via Attenuation of the RANKL/OPG Ratio. *Article.Int. J. Mol. Sci. 2014, 15, 17130-17147* ISSN 1422-0067.

¹⁸Mar'atirrosyidah R., Estiasih, T., (2015). Aktivitas Antioksidan Senyawa Bioaktif Umbi-Umbi Lokal Inferior: Kajian Pustaka. *Jurnal Pangandan Agroindustri Vol. 3 No. 2 p.594-601, April 2015.*

The Effect of Ethanolic Extract of Purple Yam Tuber (*Dioscorea alata* L.) on Bone Calcium Levels in Ovariectomized Rat

ORIGINALITY REPORT

11 %	%	11 %	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

- 1** Marie Claire Mukakarangwa, Geldine Chironda, Busisiwe Bhengu, Godfrey Katende. **2%**
"Adherence to Hemodialysis and Associated Factors among End Stage Renal Disease Patients at Selected Nephrology Units in Rwanda: A Descriptive Cross-Sectional Study", *Nursing Research and Practice*, 2018
Publication
- 2** Swamy, M, W Norlina, W Azman, D Suhaili, KNS Sirajudeen, Z Mustapha, and C Govindasamy. **1%**
"Restoration Of Glutamine Synthetase Activity, Nitric Oxide Levels And Amelioration Of Oxidative Stress By Propolis In Kainic Acid Mediated Excitotoxicity", *African Journal of Traditional Complementary and Alternative Medicines*, 2014.
Publication
- 3** Titi Komalasari, Sabtanti Harimurti. **1%**
"A Review of The Anti-diabetic Activity of *Andrographis paniculata* (Burm. f.) Nees based in-vivo Study",

4

M.H Veerendra Kumar, Y.K Gupta. "Effect of different extracts of *Centella asiatica* on cognition and markers of oxidative stress in rats", *Journal of Ethnopharmacology*, 2002

Publication

1%

5

J. Shi, Z. Liu, T. Kawai, Y. Zhou, X. Han. "Antibiotic administration alleviates the aggravating effect of orthodontic force on ligature-induced experimental periodontitis bone loss in mice", *Journal of Periodontal Research*, 2017

Publication

1%

6

Arief Nurrochmad, Supardjan Amir Margono, Sardjiman, Arief Rahman Hakim, Ernawati, Erna Kurniawati, Erva Fatmawati. "Hepatoprotective and antioxidant activity of pentagamavunon-0 against carbon tetrachloride-induced hepatic injury in rats", *Asian Pacific Journal of Tropical Medicine*, 2013

Publication

1%

7

O. Ortmann, J. M. Weiss, K. Diedrich. "Nutzen und Risiken der Hormonersatztherapie im Klimakterium und in der Postmenopause", *Gynäkologische Endokrinologie*, 2003

Publication

1%

8

Lindgren, P.R.. "Estrogen and progesterone receptors in ovarian epithelial tumors",
Molecular and Cellular Endocrinology,
20040630

Publication

1%

9

Jintanaporn Wattanathorn, Supannika
Kawvised, Wipawee Thukham-mee.
"Encapsulated Mulberry Fruit Extract Alleviates
Changes in an Animal Model of Menopause with
Metabolic Syndrome", Oxidative Medicine and
Cellular Longevity, 2019

Publication

1%

10

Cheng-Hsin Wang, Chung-Hung Tsai, Hui-Ju
Lin, Tse-Cheng Wang, Hsiao-Ling Chen.
"Uncooked Taiwanese yam (*Dioscorea alata* L.
cv. Tainung No. 2) beneficially modulated the
large bowel function and faecal microflora in
BALB/c mice", Journal of the Science of Food
and Agriculture, 2007

Publication

1%

11

J. Lu, R. N. S. Wong, L. Zhang, R. Y. L. Wong,
T. B. Ng, K. F. Lee, Y. B. Zhang, L. X. Lao, J. Y.
Liu, S. C. W. Sze. "Comparative Analysis of
Proteins with Stimulating Activity on Ovarian
Estradiol Biosynthesis from Four Different
Dioscorea Species in vitro Using Both
Phenotypic and Target-based Approaches:

1%

Implication for Treating Menopause", Applied Biochemistry and Biotechnology, 2016

Publication

Exclude quotes On

Exclude matches < 1%

Exclude bibliography Off