

Lampiran 1. Analisis Data

Crosstabs

Hemodialisa * Kreatinin Crosstabulation

| | | | Kreatinin | | Total |
|-------------|--------|----------------------|---------------|--------------|--------|
| | | | Kreatinin <=3 | Kreatinin >3 | |
| Hemodialisa | 2 kali | Count | 3 | 15 | 18 |
| | | % within Hemodialisa | 16,7% | 83,3% | 100,0% |
| | | % within Kreatinin | 27,3% | 36,6% | 34,6% |
| | | % of Total | 5,8% | 28,8% | 34,6% |
| | 3 kali | Count | 8 | 26 | 34 |
| | | % within Hemodialisa | 23,5% | 76,5% | 100,0% |
| | | % within Kreatinin | 72,7% | 63,4% | 65,4% |
| | | % of Total | 15,4% | 50,0% | 65,4% |
| Total | | Count | 11 | 41 | 52 |
| | | % within Hemodialisa | 21,2% | 78,8% | 100,0% |
| | | % within Kreatinin | 100,0% | 100,0% | 100,0% |
| | | % of Total | 21,2% | 78,8% | 100,0% |

Chi-Square Tests

| | Value | Df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|-------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | ,332 ^a | 1 | ,564 | | |
| Continuity Correction ^b | ,048 | 1 | ,826 | | |
| Likelihood Ratio | ,342 | 1 | ,559 | | |
| Fisher's Exact Test | | | | ,727 | ,422 |
| Linear-by-Linear Association | ,326 | 1 | ,568 | | |
| N of Valid Cases | 52 | | | | |

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 3,81.

b. Computed only for a 2x2 table