

DAFTAR PUSTAKA

- Abraham, S., Venu, A., Ramachandran, A., Chandran, P.M., Raman, S. (2012). Assasment of Quality of Life in Patients on Hemodialysis and the Impact of Counseling. *Saudi Journal of Kidney Disease and Transplantation*;23(5):953-957.
- Al-Arabi, S. (2006). Quality of life subjectives descriptions of challenges to patients with end stage renal disease. *Nephrol Nurs J*.33(3):285-92.
- Amro A, Waldum B, Dammen T. (2014). Symptom clusters in patients on dialysis and their association with quality-of-life outcomes. *Journal of Renal Care*;40(1):23-33.
- Anees, M., Hameed, F., Mumtaz, A., Ibrahim, M., Khan, M.N.S. (2011). Dialysis-Related Factors Affecting Quality of Life in Patients on Hemodialysis. *Iranian Journal of Kidney Disease*. Volume 5.Number 1. January.
- Anees, M., Malik., M., Abbasi, T., Nasir, Z., et al. (2014). Demographic factors affecting quality of life of hemodialysis patients. *Pak J Med Sci*.Vol.30.No.5
- Anu, VK., Pushpa, P., Kumar, S. (2013). Quality of life of patients undergoing haemodialysis at B.P.Koirala insitiute of Health Sciences. ISSN:2091-1041.Volume 1.Issue 2.
- Apostolou, T, Hutchison, AJ, Boulton AJ, Chak W, Vileikyte L. (2007). Quality of life in CAPD, transplant, and chronic renal failure patients with diabetes. *Ren.Fail*,29 (2):189-187.
- Arikunto, S. (2006). *Prosedur Penelitian Suatu Pendekatan Praktik*.Ed.Revisi VI.PT Rineka Cipta. Jakarta.
- Asri, P, Marthon, Marjono, Purwanto. (2006). Hubungan dukungan sosial dengan tingkat depresi pasien yang menjalani terapi hemodialisa. Mei 23,2016. <http://www.i-lib.ugm.ac.id>
- Azwar, S. (2005). *Sikap Manusia:Teori dan Pengukurannya*. Pustaka Pelajar. Yogyakarta.
- Bakewell, A.B., Higgins, R.M., & Edmunds, M.E. (2002). Quality of life in peritoneal dialysis patients:Decline over time and association with clinical outcomes. *Kidney International*.61,239-248.

- Beddhu, S., Brun, F.J., Saul, M., Seddon, P., et al. (2000). A simple comorbidity scale predicts clinical outcome and cost in dialysis patients. *The American Journal of Medicine*, 108, 609-613.
- Black, J.M., & Hawk, J.H. (2005). *Medical Surgical Nursing : Clinical Management for Positive Outcome*. 7th edition. Philadelphia : W.B. Saunders Company.
- Braga, S.F., Peixoto, S.V., Gomes, I.C., Acurcio, F., et al. (2011). Factors associated with health-related quality of life in elderly patients on hemodialysis. *Rev Saude Publica*; 45 (6).
- Brenner, B.M. (2004). *Brenner & Rector's. The Kidney*. 7th. Edition. Philadelphia: Elsevier.
- Cavanaugh, K.L. (2007). Diabetes Management Issue for Patients With Chronic Kidney Disease. *Clinical Diabetes*. Volume 25, Number 3.
- Cella, D.F. (1992). Method and problems in measuring quality of life. *Support Care Cancer*. 3:11-22.
- Cianci, R., S.Lai, L.Fuiano, A. Gigante., P. (2009). Hypertension in Hemodialysis. An Overview on Physiopathology and Therapeutic Approach in Adults and Children. *The Open Urology & Nephrology Journal*, 2 : 11-19.
- Coccosis, M.G., Theofilou, P., Synodinou, C., Tomars, V., et al. (2008). Quality of life, mental health and health beliefs in haemodialysis and peritoneal dialysis patients : Investigating differences in early and later years of current treatment. *BMC Nephrology*, 9:14.doi:10.1186/1471-2369-9-14.
<http://www.biomedcentral.com/1471-2369/9/14>
- Eckersten, D. (2015). Anti-Mullerian hormone. A Sertoli cell-derived marker, is decreased in plasma of male patients in all stages of chronic kidney disease. *Andrology*. 3(6):1160-4.
- El-Shaded, A.M., Sharf, S.A., El-Sebae, H.A., Roshdy, M.M. (2013). Hemoglobin Level, Associated Co-Morbidities and Quality of Life among Patients Undergoing Hemodialysis at One of the University Hospitals in Cairo Governorate. *World Applied Sciences Journal*. 23(1):29-36.
- Evans, M. (2005). The natural history of chronic renal failure: results from an unselected, population-based, inception cohort in Sweden. *Am J of Kidney Dis*. 46(5):863-70.

- Farquahar, M. (1995). Elderly people definition of quality of life, social science and medicine, 41, 1436-1446.
- Fried, T.R., O'Leary, J., Towle, V., Goldstein, M.K., et al. (2014). The Effects of Comorbidity on the Benefits and Harms of Treatment for Chronic Disease : A Systematic Review. *PLoS ONE* 9(11) : e112593. doi:10.1371/journal.pone.0112593
- Ghaderian S.B., Beladi-Mosavi S.S. (2015). The role of diabetes and hypertension in chronic-kidney disease.*J Renal Inj Prev.*3(4):109-110.
- Gerasimoula, K., Lefkothea, L., Maria, L., Victoria, A., et al. (2015). Quality of Life in Hemodialysis Patients. *Mater Sociomed.*Oct; 27 (5); 305-309.
- Goldberg I., Krause I. (2016). The Role of Gender in Chronic Kidney Disease. *EMJ.*2016:1(2):58-64.
- Gopalan. (2008). *Eight-point Risk Factor Checklist Evolved to Predict Chronic Kidney Disease*. Available from : <http://www.medindia.net/mews/Eight-point-risk-factor-checklist-evolved-to-predict-chronic-kidney-disease-45537-2-htm> (Accessed 20 Februari 2016).
- Guerrero, V., Alvarado, O., Espina, M. (2012). Quality of life in people with chronic hemodialysis:association with sociodemographic, medical-clinical and laboratory variables.*Rev.Latino-Am.Enfermagen.*Sept-Oct;20(5):838-46.
- Gumprecht, J., Zelobowska, K., Gosek, K., Zywiec, J., et al. (2010).Quality of life among diabetic and non-diabetic patients on maintenance haemodialysis. *Exp. Clin. Endocrinol Diabetes.*118(3):205-208.
- Gunal, A.I., Duman, S., Ozkahya. (2001). Strict volume control normalizes hypertension in peritoneal dialysis patients. *American Journal of Kidney Disease.* Vol 37, No.3, pp 588-593.
- Haddiya, I., R. El Harraqui, I., Karimi, N. Benabdallah and Y. Bentata. (2013). Therapeutic Education's Role in the Management of Hypertension in Chronic Hemodialysis Patients. *ISRN Hypertension*, pp:1-7
- Hastono, S.P. (2007). *Analisis Data Kesehatan*. Fakultas Kesehatan Masyarakat. Universitas Indonesia: Jakarta.

- Ignatavicius, D.D., & Workman, M.L. (2006). *Medical Surgical Nursing: Critical Thinking for Collaborative Care*. 5th edition. St Louis: Elsevier Saunders.
- Imai E., Horio M., Watanabe, T., Iseki, K., et al. (2009). Prevalence of chronic kidney disease in the Japanese general population. *Clinical and Experimental Nephrology*. December 2009, 13:621.
- Istanti, Y.P. (2009). Faktor-faktor yang berkontribusi terhadap interdialytic weight gain (IDWG) di Unit Hemodialisa RS PKU Muhammadiyah Yogyakarta. *Jurnal Universitas Indonesia*.
- Indonesian Renal Registry. (2010). 4th Report of Indonesian Renal Registry. IRR: 18-20.
- Jha, V. (2004). End-stage renal care in developing countries. The India Experience. *Ren Fail.*;26:201-8.
- Jofre, R., Juan, M., Lopez-Gomez, Valderrabano, F. (2000). Quality of life for patient groups. *Kidney International*, Vol.57, Suppl,74, pp S-121-S-130.
- Jonge, P., Ruinemans, G., Huyse, F., Wee, P. (2003). A simple risk score predicts poor quality of life and non-survival at 1 year follow-up in dialysis patients. *Nephrol Dial Transplant*.18: 2622-2628. doi:10.1093/ndt/gfg453.
- Juergesen E., Wuerth D, Finkelstein SH, Juergersen PH., et al. (2006). Hemodialysis and peritoneal dialysis: Patients' assessment of their satisfaction with therapy and the impact of the therapy on their lives. *J Am Soc Nephrol*;1:1191-6.
- Kan, W-C., Wang, J-J., Wang, S-Y., Sun, Y-M., et al. (2013). The New Comorbidity Index for Predicting Survival in Elderly Dialysis Patients : A Long-Term Population-Based Study. *PLUS ONE*, 8(8), e68748
- Khaled A.K., Larisa M., Karpov, I. (2009). Individual quality of life in chronic kidney disease: Influence of age and dialysis modality. *Clin J Ann Soe Nephrol*;4:711-8.
- Khaw, W.F., Hassan, S.T., Latiffah, A.L. (2011). Health-related Quality of Life among Hypertensive Patients Compared with General Population Norms. *Journal of Medical Sciences*, 11:84-89.
- Kinghorn, S., Gamlin, R. (2004). *Palliative Nursing: Bringing Comfort and Hope*, Bailliere, Tindall, St. Louise.

- Kiran, V.U., N.V. Rajaiah, D.R. Krishna and Y.N. Reddy. (2010). Effect of losartan and ramipil on oxidative stress and anti-oxidant status in South Indian hypertensive patients. *Int.J.Pharmacol*, 6: 916-920.
- Kummer, S. (2012). The influence of gender and sexual hormones on incidence and outcome of chronic kidney disease. *Pediatr Nephrol*.27(8):1213-9.
- Kring, D.L., & Crame, P.B. (2009). Factors affecting quality of life in persons on hemodialysis. *Nephrology Nursing Journal*. 36, 15-24.
- Krylova MI, Shutov EV, Ermolenko VM. (2010). Survival and Quality of Life of Patients with Renal Replacement Therapy. *Vestnik YGU*;2:63-70.
- Lameire, N., Mehta, R.L. (2000). *Complications of dialysis*. New York: Informa Health Care.
- Levin. (2008). *Guidelines for the management chronic kidney disease*. January 5, 2015. <http://www.cmaj.ca/>
- Le Mone, P., & Burke, K.M. (1996). *Critical Thinking in Client Care*. Addison Wesley.
- Le Mone, P., & Burke, K.M. (2008). *Medical Surgical Nursing : Critical Thinking in Client Care*, 6th edition, New Jersey : Prentice Hall Health.
- Liu, W.J., Chew, T.F., Chiu, A.S., Zaki, M. (2006). Quality of life of dialysis patients in Malaysia, *Med J Malaysia*;61 (513-5).
- Mandoorah, Q.M., Shaheen, F.A., Mandoorah, S.M., Bawazir, S.A., et al. (2014). Impact of Demographic and Comorbid Conditions on Quality of Life of Hemodialysis Patients: A Cross-sectional Study. *Saudy J Kidney Dis Transpl*;25 (2):432-437.
- Makkar, V., Kumar, M., Mahajan, R., Khaira, N.S. (2015). Comparasion of Outcomes and Quality of Life between Hemodialysis and Peritoneal Dialysis Patients in India ESRD Population. *Journal of Clinical and Diagnostic Research*.Mar, Vol -9(3): OC28-OC31.
- Malliara, M. (2007). The Management of Hypertension in hemodialysis and CAPD patients. *Hippokratia*, 11 (4): 171-174.
- Mansjoer, A., Triyanti,, K., Savitri, R., Wardhani, W.I. (2001). *Kapita Selekt Kedokteran*. Ed.3. Media Aesculapis, Jakarta.

- McCullough, P.A. (2004). Cardiovascular disease in chronic kidney disease from cardiologist's perspective. *Curr Opin Nephrol Hypertens*;13:591-600.
- McIntyre, C.W. (2010). Haemodialysis-induced myocardial stunning in chronic kidney disease-a new aspect of cardiovascular disease. *Blood Purif*;29:105-110.
- Moltó, A., Dougados, M. (2014). Comorbidity indices. *Clin Exp Rheumatol*;32 (Suppl.85): S131-S134.
- Morton, R.L. Snelling, P., Webster, A.C., Rose, J., et al. (2012). Factor influencing patient choice of dialysis versus conservative care to treat end-stage kidney disease. *CMAJ*. March 20, 184 (5).
- Murphy, B. (2000). Australian WHOQOL-100, WHOQOL-BREF and CA-WHOQOL-INSTRUMENTS:user manual and interpretation guide. February 23, 2016. <http://www.psychiatry.unimelb.edu.au/>
- National Kidney Foundation. (2009). Chronic Kidney Disease. New York: National Kidney Foundation. Available from: <http://www.kidney.org/kidneydisease/ckd/index.cfm#whatits>.
- Neugarten J., Acharya, A., Silbiger, S.R. (2000). Effect of gender on the progression of nondiabetic renal disease a meta-analysis. *J Am Soc Nephrol*;11(2):319-29.
- Nurchayati, S. (2011). *Analisis faktor-faktor yang berhubungan dengan kualitas hidup pasien penyakit ginjal kronik yang menjalankan hemodialisa di Rumah Sakit Islam Fatimah Cilacap dan Rumah Sakit Umum Daerah Banyumas*. Tesis. Universitas Indonesia.
- Pakpour, A., Saffari, M., Yekaninejad, S., Panahi, D., et al. (2010). Health-Related Quality of Life in a Sample of Iranian Patients in Hemodialysis. *Iranian Journal of Kidney Disease*. Volume 4. Number 1. January.
- Paraskevi, T. (2011). The role of sociodemographic factors in health-related quality of life of patients with end-stage renal disease. *International Journal of Caring Sciences*. January-April. Vol 4. Issue 1.
- Paraskevi, T. (2012). The Impact of Sociodemographic and Psychological variables on quality of life in patients with renal disease : Findings of a cross-sectional study in Greece. *World J Nephrol Urol*:1 (4-5):101-106.
- PERNEFRI. (2012). *4th Report Of Indonesian Renal Registry*.

- Rirava MB, Robinson-Cohen C, Kestenbaum B. (2015). Changes in symptom burden and physical performance with initiation of dialysis in patients with chronic kidney disease. *Hemodialysis International*:19:147-150.
- Saad.E., Charra, B., Raj, D.SC. (2004). Hypertension control with daily dialysis. *Seminars in Dialysis*. Vol 17, N.4. pp 295-298.
- Saboya, P.,P.R. Zimmermann and L.C. Bodanese. (2010). Association between anxiety or depressive symptoms and arterial hypertension and their impact on the quality of life. *Int.J.Psychiatry Med.*, 40: 307-320.
- Salyzhyn, T.I., Yatsyshym, R.I. (2015). Quality of Life of Patients with Chronic Kidney Disease Receiving Hemodialysis with Concomitant Chronic Heart Failure and without it. *Galician Medical Journal*;22 (4)
- Sarwono, J. (2006). *Metode Penelitian Kuantitatif dan Kualitatif*. Graha Ilmu.Yogyakarta.
- Sathvik, B.S., Parthasarathi, G., Narahari, M.G., Gurudev, K.C. (2010). An assessment of the quality of life in hemodialysis patients using the WHOQOL-BREF questionnaire. *Indian Journal of Nephrology*.October.Vol 18.Issue 4.
- Shahgholian, N., Ghafourifard, M., Rafieian, M., & Mortazavi, M. (2008). Impact of two types of sodium and ultrafiltration profiles on intradialytic hypotension in hemodialysis patients. *IJNMR Autumn*.13, 135-136.
- Shiba, N., Shimokawa, H. (2011).Chronic Kidney disease and heart failure-Bidirectional close link and common therapeutic goal. *Journal of Cardiology*.57, 8-17.
- Smeltzer, S. C & Bare B.G. (2008). *Brunner & Suddarth's Textbook of medical-surgical nursing* (11th Ed). Philadelphia: Lipponcott William& Wilkins.
- Sorensen, V.R., Mathiesen, R. Watt, T., Bjorner, J.B., et al. (2007). Diabetic patients treated with dialysis:complications and quality of life. *Diabetologia*:50:2254-2262.
- Sugiharto. (2007). *Faktor-faktor risiko hipertensi grade II pada masyarakat*. Disertasi. Universitas Diponegoro Semarang.
- Sugiyono. (2005). *Statistik untuk Penelitian*. PT Rineka Cipta: Jakarta.

- Suwitra, K. (2006). *Penyakit Ginjal Kronik*, dalam Sudoyono, A.W.Sutiyahadi, B., Alwi, L., Simadibrata, M., & Setiati, S. Buku Ajar Ilmu penyakit dalam (581) Jakarta : Departemen Ilmu Penyakit Dalam FK UI.
- Tedla, E.M., Baar, A., Browne, R., Brown, C. (2011).Hypertension in Chronic Kidney Disease :Navigating the Evidence. *International Journal of Hypertension*.Article ID 132405, 9 pages.
- The Kidney Disease of Diabetes page. National Institute of Diabetes & Digestive & Kidney Disease. Web site. Available at: <http://www.niddk.nih.gov/health/kidney/pubs/kdd/kdd/htm>. Diunduh pada tanggal 30 Mei 2016.
- Tisher, C.C., Wilcox, C.S. (1995). *Nefrologi*. EGC: Jakarta.
- UKPDS. (1998). Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes:UKPDS 38.UK Prospective Diabetes Study Group.*BMJ*;317:703-13.
- Ulya,I., Suryanto. (2005). Perbedaan Kadar Hb Pre dan Post Hemodialsa pada Penderita Gagal Ginjal Kronis di RS PKU Muhammadiyah Yogyakarta.*Mutiara Medika*.Vol 7, No.1:29-33.
- Unruh, M.L., Newman, A.B., Larive, B., Dew, M.A., et al. (2008). The influence of age on changes in helath-related quality of life over three years in a cohort undergoing hemodialysis. *J Am Geriatr Soc*.56(9):1608-17.
- Van, K.N., Duangpaeng, S., Deenan, A., & Bonner, A. (2012). Examining the health-related quality of life of people with end-stage kidney disease living in Hanoi, Vietnam. *Renal Society of Australasia Journal*, 8(3), 140-145.
- Wahid, A., Bokhari, S.A.H., Butt, S., Khan, A.A. (2014). Quality of life in diabetic and non diabetic patients on hemodialysis therapy. *Journal of Diabetes and Endocrinology*.Vol 5(2),pp.9-18, February.
- White C, McDonnel H. (2014). Psychosocial distress in patients with end-stage kidney disease. *Journal of Renal Care*:40:78-81.
- World Health Organization. (1996). *WHOQOL-BREF:Introduction, administration, scoring and generic version of assessment*.Geneva:WHO.

Yang S.C, Kuo P.W, Wang J.D, Lin Mi, Su S. (2005). Quality of life and its determinants of hemodialysis patients in Taiwan measured with WHOQOL-BREF (TW). *Am.J.Kidney Dis.*46(4):635-641.

Yuwono. (2000). *Kualitas Hidup menurut Spitzer pada Penderita Gagal Ginjal Terminal yang Menjalankan Hemodialisa di Unit Hemodialisa RSUP. Kariadi, Semarang.*

Zadeh, K.K. (2003). *Quality of Life in Patients with Chronic Renal Failure*. February 20, 2016. <http://www.kidney.org>